

ADHD-LIKE SYMPTOMS IN CHILDREN AFFECTED BY OBSTRUCTIVE SLEEP APNEA SYNDROME: A CASE-CONTROL STUDY

FRANCESCO PRECENZANO*, MARIA RUBERTO**, LUCIA PARISI***, MARGHERITA SALERNO****, AGATA MALTESE***, ILARIA D'ALESSANDRO*, IMMACOLATA DELLA VALLE*, GENNARO VISCO*, ROSARIA MARTINA MAGLIULO*, GIOVANNI MESSINA*****, MICHELE ROCCELLA***

*Clinic of Child and Adolescent Neuropsychiatry; Headache Center for children and adolescents, Department of Mental Health and Physical and Preventive Medicine, Second University of Naples - **Department of Medical-Surgical and Dental Specialties, Second University of Naples - ***Department of Psychological, Pedagogical and Educational Sciences, University of Palermo, Italy - ****Sciences for Mother and Child Health Promotion, University of Palermo, Italy - *****Department of Experimental Medicine, Section of Human Physiology and Unit of Dietetics and Sports Medicine, Department of Clinical and Experimental Medicine, University of Foggia, Foggia, Italy

#Francesco Precenzano and Maria Ruberto equally contributed to the manuscript

ABSTRACT

Introduction: ADHD is characterized by inattention, hyperactivity, impulsivity, or a combination of these symptoms. Sleep disorders may be considered as a not secondary underlying cause of ADHD and growing evidence evidenced that obstructive sleep apnea syndrome (OSA) symptoms may overlap ADHD's ones.

Aim of the present study is verifying the presence of ADHD-like symptoms in other frequent condition such as sleep-related breathing disorders in pediatric age.

Materials and methods: 34 children (19 males and 15 females) aged 6-10 years (mean age 9.706 ± 3.434) with polysomnographic diagnosis of OSA, according to ICSD-3 criteria, were recruited. Control group was composed by 89 typical developing children (47 males, 42 females) (mean age 9.528 ± 3.351). The Conners' Rating Scale- Revised (CRS-R) test was administered to mothers to assess their children's behavior.

Results: The two groups are comparable for age ($p = 0.794$) and sex distribution ($p = 0.918$). Subjects affected by OSA present higher scores in pathological range in all scales of CRS-R than controls (Table 1).

Conclusions: Scientific and clinical evidences tend to stress the similarities between ADHD and OSA effects/comorbidities in pediatric age, suggesting sleep screening as mandatory before starting stimulant drugs treatment.

Keywords: ADHD, ADHD-like symptoms, obstructive sleep apnea syndrome, OSA.

DOI: 10.19193/0393-6384_2016_6_159

Received May 30, 2016; Accepted September 02, 2016

Introduction

In developmental age, the syndrome named as Attention Deficit Hyperactivity Disorder (ADHD) is characterized by inattention, hyperactivity, impulsivity, or a combination of these symptoms, compromising basic everyday functions such as learning abilities, executive functioning, and social skills⁽¹⁾.

Since the first description in 1902 by George Still on Lancet journal, the prevalence studies have

reported different prevalence rates ranging 3% -5% of school-age children, and around 1% in Italian pediatric population⁽²⁾.

To date, diagnostic criteria are based focus on behavioral symptoms, because the absence of specific biomarkers, although genetic factors, environmental factors, prematurity, low birth weight and hypoxic-ischemic encephalopathy are considered important in ADHD etiology. In this light, the role of sleep disorders may be an interesting and to date not yet well considered as causative factor for

ADHD, despite of many and relevant scientific evidences⁽⁹⁾. In fact, sleep disorders may be considered as a not secondary underlying cause of ADHD⁽³⁾, although still underdiagnosed in pediatric age. On the other hand, there is growing evidence that sleep disorders and mainly the obstructive sleep apnea syndrome (OSA) are associated with behavioral problems and hyperactivity, executive functions impairment, cognitive and learning difficulties⁽⁴⁻¹³⁾.

Starting from the frequent over-diagnosis among Italian children, the aim of the present study is verifying the presence of ADHD-like symptoms in other frequent condition such as sleep-related breathing disorders in pediatric age.

Material and methods

34 children (19 males and 15 females) aged 6-10 years (mean age 9.706 ± 3.434) with polysomnographic diagnosis of OSA, according to ICSD-3 criteria (14), were recruited.

Exclusion criteria were the following: overweight, obesity, cognitive disability (IQ <70), borderline intellectual functioning, neurological and psychiatric disorders (15-17), chromosomal defects, specific neuropsychological disorder, epilepsy⁽¹⁸⁻³¹⁾.

Control group was composed by 89 typical developing children (47 males, 42 females) (mean age $9,528 \pm 3,351$).

All subjects of both groups were recruited within the same urban area, all were Caucasian and homogeneous for socioeconomic status

Ethical approval from the local Research Ethics Center and informed parental consent were obtained.

Polysomnographic (PSG) Data

After reviewing and analyzing PSG data collected from inpatients children between January and June 2016 to establish the presence of OSA in the experimental group, OSA severity was determined according to the current guidelines specified by the American Academy of Sleep Medicine (AASM)⁽³²⁾: mild OSA was defined by an obstructive apnea-hypopnea index (o AHI) of 1 to <5 events per hour; moderate OSA was defined as was defined as ≥ 5 to <10 events/hour, and severe OSA ≥ 10 events/hour.

Conners' Rating Scale-Revised (CRS-R)

At the clinic visit, Conners' Rating Scale-Revised (CRS-R) was administered to mothers to

assess their children's behavior. CRS-R examines seven types of behavior problems in children including: Oppositional, Anxious-Shy, Cognitive Problem/Inattention, Hyperactivity, Perfectionism, Psychosomatic, and Social Problems. It also contains four index scores: Conners' ADHD Index; Conners' Global Index (CGI): Restless-Impulsive; CGI: Emotional Lability; and CGI: Total. CRS-R also includes scales for the Diagnostic and Statistical Manual of Mental Disorders, 4th Edition⁽³³⁾ diagnosis of ADHD:

- a) DSM-IV: Inattentive,
- b) DSM-IV: Hyperactive-Impulsive,
- c) DSM IV: Total.

Higher scores indicate an increased likelihood that a child would meet diagnostic criteria for ADHD. A T-score of 65 indicates a "clinically significant problem"⁽³⁴⁾.

Statistical analysis

For comparison between the two groups (OSAS and controls) it was applied t- testing and Chi-square test, where appropriate. P values <0.05 were considered statistically significant.

For statistical analysis it used the software STATISTICA (data analysis software system, version 6, StatSoft, Inc. (2001). For the comparison between the groups the t-test was applied. P values <0.05 were considered statistically significant.

Results

The two groups are comparable for age ($p = 0.794$) and sex distribution ($p = 0.918$). Subjects affected by OSA present higher scores in pathological range than controls in the following domains: hyperactivity, ADHD Index, Restlessness-impulsivity (CGI II), emotional instability (CGI IE), Hyperactivity and other problem areas (CGI T), DSM-IV inattention, DSM -IV hyperactivity and impulsivity, total DSM-IV ADHD, No. Inattention symptoms of DSM-IV, no symptoms of hyperactivity / impulsivity DSM-IV ($p < 0.001$ for all subscales) (Table 1).

Discussion

Several studies have focused on alterations of executive functioning in children with OSA, with a putative specific relationship with the degree of respiratory troubles, supporting the role of intermittent hypoxia impact the fronto-prefrontal regions^(6,7,10,35).

Considering the more relevant and frequency of OSA respect of ADHD (27% vs. 3-5%) in pediatric age, it is obvious and intuitive that hyperactive school-aged children may be affected by OSA more probable than ADHD.

	OSA N=34	Controls N=89	P
Oppositional	51.02±8.14	49.96±9.03	0.551
Cognitive Problem/Inattention	45.13±6.12	43.01±8.05	0.168
Hyperactivity	71.31±4.18	44.15±6.34	<0.001
Anxious-Shy	40.73±2.17	39.96±2.83	0.155
Perfectionism	46.93±9.15	44.18±8.93	0.132
Social Problems	50.37±7.39	50.93±7.25	0.704
Psychosomatic Problems	50.15±5.03	49.82±5.44	0.759
Conners' ADHD Index	77.36±10.04	45.26±6.19	<0.001
Restless-Impulsive (CGI I-I)	71.86±9.93	44.64±5.61	<0.001
Emotional Lability (CGI I-E)	59.05±6.94	45.11±7.09	<0.001
Hyperactivity and other problems (CGI T)	70.48±9.81	44.25±6.33	<0.001
DSM-IV Inattention	70.66±8.96	49.29±5.90	<0.001
DSM-IV Hyperactivity	75.71±8.83	49.63±4.31	<0.001
DSM-IV ADHD Total score	75.04±10.25	48.06±6.38	<0.001

Table 1: Shows differences between children affected by obstructive sleep apnea syndrome (OSA) and typical developing subjects among Conners Rating Scales-Revised version (CRS-R) scales. For comparison between the two groups (OSAS and controls) it was applied t- testing and Chi-square test, where appropriate. *P* values<0.05 were considered statistically significant.

In general, hyperactivity presents higher prevalence as non-specific symptom in several nosographic paintings of childhood neuropsychiatric interest, as a result of a reduction in inhibitory control both motor and behavioral skills. In fact, hyperactive and inattentive behavior has been reported in children with OSA and habitual snoring. In addition, children with higher indices of snoring tend to show lower scores to cognitive and neuropsychological standardized tests. On the other hand, in 2011 a systematic review has shown that the prevalence of OSA in ADHD subjects (25-30%) is higher than that of the general population (about 3%)⁽³⁶⁾. In this perspective, the present study has as preliminary investigation in the evaluation of hyperactive behavior in school-aged children with OSAS.

Conversely, sleepy children tend to use a greater motor activity pattern to be more vigilant, supporting the parents' report about that when they are very tired appear angry and become aggressive and hyperactive. In this light, the use of stimulant drugs increase the catecholamines activity in the CNS, improving arousal and alertness in typical developing children, identifying it as the best treatment in ADHD to limit motor hyperactivity⁽²⁷⁻²⁹⁾.

Moreover, the hypothesis that OSA may represent one of the causes of ADHD is supported by AASM guidelines that recommend screening for sleep apnea among children undergoing evaluation for ADHD⁽⁴⁰⁾, because of the overlapping cognitive, neuropsychological and behavioral symptoms and OSA.

In conclusion, scientific and clinical evidences tend to stress the similarities between ADHD and OSA effects/comorbidities in pediatric age, suggesting sleep screening as mandatory before starting stimulant drugs treatment⁽⁴¹⁻⁵⁰⁾.

References

- 1) Craig F, Margari F, Legrottaglie AR, Palumbi R, de Giambattista C, Margari L. *A review of executive function deficits in autism spectrum disorder and attention-deficit/hyperactivity disorder*. Neuropsychiatr Dis Treat. 2016 May 12; 12: 1191-202. doi: 10.2147/NDT.S104620
- 2) Cortese S, Pani P, Arcieri R, Germinario EA, Capuano A, Margari L, Chiarotti F, Curatolo P. *Safety of Methylphenidate and Atomoxetine in Children with Attention-Deficit/Hyperactivity Disorder (ADHD): Data from the Italian National ADHD Registry*. CNS Drugs. 2015; 29(10): 865-77. doi: 10.1007/s40263-015-0266-7
- 3) Miano S, Esposito M, Foderaro G, Ramelli GP, Pezzoli V, Manconi M. *Sleep-Related Disorders in Children with Attention-Deficit Hyperactivity Disorder: Preliminary Results of a Full Sleep Assessment Study*. CNS Neurosci Ther. 2016 Jun 3. doi: 10.1111/cns.12573
- 4) Santamaria F, Esposito M, Montella S, Cantone E, Mollica C, De Stefano S, Mirra V, Carotenuto M. *Sleep disordered breathing and airway disease in primary ciliary dyskinesia*. Respirology. 2014 May; 19(4): 570-5. doi: 10.1111/resp.12273
- 5) Carotenuto M, Esposito M, Di Pasquale F, De Stefano S, Santamaria F. *Psychological, cognitive and maternal stress assessment in children with primary ciliary dyskinesia*. World J Pediatr. 2013 Nov; 9(4): 312-7. doi:10.1007/s12519-013-0441-1
- 6) Esposito M, Antinolfi L, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Precenzano F, Carotenuto M. *Executive dysfunction in children affected by obstructive sleep apnea syndrome: an observational study*. Neuropsychiatr Dis Treat. 2013; 9: 1087-94. doi: 10.2147/NDT.S47287

- 7) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Patriciello G, Precenzano F, Carotenuto M. *Visuomotor competencies and primary monosymptomatic nocturnal enuresis in prepubertal aged children*. *Neuropsychiatr Dis Treat*. 2013; 9: 921-6. doi: 10.2147/NDT.S46772.
- 8) Carotenuto M, Gimigliano F, Fiordelisi G, Ruberto M, Esposito M. *Positional abnormalities during sleep in children affected by obstructive sleep apnea syndrome: the putative role of kinetic muscular chains*. *Med Hypotheses*. 2013 Aug; 81(2): 306-8. doi: 10.1016/j.mehy.2013.04.023.
- 9) Esposito M, Gallai B, Parisi L, Roccella M, Marotta R, Lavano SM, Mazzotta G, Carotenuto M. *Primary nocturnal enuresis as a risk factor for sleep disorders: an observational questionnaire-based multicenter study*. *Neuropsychiatr Dis Treat*. 2013; 9: 437-43. doi: 10.2147/NDT.S43673.
- 10) Carotenuto M, Esposito M, Parisi L, Gallai B, Marotta R, Pascotto A, Roccella M. *Depressive symptoms and childhood sleep apnea syndrome*. *Neuropsychiatr Dis Treat*. 2012; 8: 369-73. doi: 10.2147/NDT.S35974.
- 11) Carotenuto M, Esposito M, D'Aniello A, Ripa CD, Precenzano F, Pascotto A, Bravaccio C, Elia M. *Polysomnographic findings in Rett syndrome: a case-control study*. *Sleep Breath*. 2013 Mar; 17(1): 93-8. doi: 10.1007/s11325-012-0654-x.
- 12) Esposito M, Carotenuto M, Roccella M. *Primary nocturnal enuresis and learning disability*. *Minerva Pediatr*. 2011 Apr; 63(2): 99-104. PubMed PMID: 21487372.
- 13) Carotenuto M, Esposito M, Pascotto A. *Facial patterns and primary nocturnal enuresis in children*. *Sleep Breath*. 2011 May; 15(2): 221-7. doi:10.1007/s11325-010-0388-6.
- 14) American Academy of Sleep Medicine. *International Classification of Sleep Disorders*. 3rd ed. Darien, IL: American Academy of Sleep Medicine; 2014.
- 15) Guzzetta A, Pizzardi A, Belmonti V, Boldrini A, Carotenuto M, D'Acunto G, Ferrari F, Fiori S, Gallo C, Ghirri P, Mercuri E, Romeo D, Roversi MF, Cioni G. *Hand movements at 3 months predict later hemiplegia in term infants with neonatal cerebral infarction*. *Dev Med Child Neurol*. 2010 Aug; 52(8): 767-72. doi:10.1111/j.1469-8749.2009.03497.x.
- 16) Esposito M, Roccella M, Parisi L, Gallai B, Carotenuto M. *Hypersomnia in children affected by migraine without aura: a questionnaire-based case-control study*. *Neuropsychiatr Dis Treat*. 2013; 9: 289-94. doi: 10.2147/NDT.S42182.
- 17) Esposito M, Ruberto M, Pascotto A, Carotenuto M. *Nutraceutical preparations in childhood migraine prophylaxis: effects on headache outcomes including disability and behaviour*. *Neurol Sci*. 2012 Dec; 33(6): 1365-8. doi: 10.1007/s10072-012-1019-8.
- 18) Carotenuto M, Santoro N, Grandone A, Santoro E, Pascotto C, Pascotto A, Perrone L, del Giudice EM. *The insulin gene variable number of tandem repeats (INS-VNTR) genotype and sleep disordered breathing in childhood obesity*. *J Endocrinol Invest*. 2009 Oct; 32(9): 752-5. doi: 10.3275/6398.
- 19) Messina A, De Fusco C, Monda V, Esposito M, Moscatelli F, Valenzano A, Carotenuto M, Viggiano E, Chieffi S, De Luca V, Cibelli G, Monda M, Messina G. *Role of the Orexin System on the Hypothalamus-Pituitary-Thyroid Axis*. *Front Neural Circuits*. 2016 Aug 25; 10: 66. doi: 10.3389/fncir.2016.00066.
- 20) Perillo L, Esposito M, Caprioglio A, Attanasio S, Santini AC, Carotenuto M. *Orthodontic treatment need for adolescents in the Campania region: the malocclusion impact on self-concept*. *Patient Prefer Adherence*. 2014 Mar 19; 8: 353-9. doi: 10.2147/PPA.S58971
- 21) Esposito M, Carotenuto M. *Borderline intellectual functioning and sleep: the role of cyclic alternating pattern*. *Neurosci Lett*. 2010 Nov 19; 485(2): 89-93. doi:10.1016/j.neulet.2010.08.062.
- 22) Esposito M, Carotenuto M. *Intellectual disabilities and power spectra analysis during sleep: a new perspective on borderline intellectual functioning*. *J Intellect Disabil Res*. 2014 May; 58(5): 421-9. doi: 10.1111/jir.12036.
- 23) Pasquali D, Carotenuto M, Leporati P, Esposito M, Antinolfi L, Esposito D, Accardo G, Carella C, Chiovato L, Rotondi M. *Maternal hypothyroidism and subsequent neuropsychological outcome of the progeny: a family portrait*. *Endocrine*. 2015 Dec; 50(3): 797-801. doi: 10.1007/s12020-015-0564-3
- 24) Franzoni E, Matricardi S, Di Pisa V, Capovilla G, Romeo A, Tozzi E, Pruna D, Salerno GG, Zamponi N, Accorsi P, Giordano L, Coppola G, Cerminara C, Curatolo P, Nicita F, Spalice A, Grosso S, Pavone P, Striano P, Parisi P, Boni A, Gobbi G, Carotenuto M, Esposito M, Cottone C, Verrotti A. *Refractory absence seizures: An Italian multicenter retrospective study*. *Eur J Paediatr Neurol*. 2015 Nov; 19(6): 660-4. doi: 10.1016/j.ejpn.2015.07.008.
- 25) Esposito M, Gallai B, Roccella M, Marotta R, Lavano F, Lavano SM, Mazzotta G, Bove D, Sorrentino M, Precenzano F, Carotenuto M. *Anxiety and depression levels in prepubertal obese children: a case-control study*. *Neuropsychiatr Dis Treat*. 2014 Oct 3; 10:1897-902. doi: 10.2147/NDT.S69795.
- 26) Esposito M, Roccella M, Gallai B, Parisi L, Lavano SM, Marotta R, Carotenuto M. *Maternal personality profile of children affected by migraine*. *Neuropsychiatr Dis Treat*. 2013; 9: 1351-8. doi: 10.2147/NDT.S51554.
- 27) Esposito M, Marotta R, Gallai B, Parisi L, Patriciello G, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. *Temperamental characteristics in childhood migraine without aura: a multicenter study*. *Neuropsychiatr Dis Treat*. 2013; 9: 1187-92. doi:10.2147/NDT.S50458.
- 28) Esposito M, Gallai B, Parisi L, Castaldo L, Marotta R, Lavano SM, Mazzotta G, Roccella M, Carotenuto M. *Self-concept evaluation and migraine without aura in childhood*. *Neuropsychiatr Dis Treat*. 2013; 9: 1061-6. doi: 10.2147/NDT.S49364.
- 29) Esposito M, Ruberto M, Pascotto A, Carotenuto M. *Nutraceutical preparations in childhood migraine prophylaxis: effects on headache outcomes including disability and behaviour*. *Neurol Sci*. 2012 Dec; 33(6): 1365-8. doi: 10.1007/s10072-012-1019-8.
- 30) Carotenuto M, Esposito M, Precenzano F, Castaldo L, Roccella M. *Cosleeping in childhood migraine*. *Minerva Pediatr*. 2011 Apr; 63(2): 105-9.
- 31) Verrotti A, Cusmai R, Laino D, Carotenuto M, Esposito M, Falsaperla R, Margari L, Rizzo R, Savasta S, Grosso S, Striano P, Belcastro V, Franzoni E, Curatolo

- P, Giordano L, Freri E, Matricardi S, Pruna D, Toldo I, Tozzi E, Lobefalo L, Operto F, Altobelli E, Chiarelli F, Spalice A. *Long-term outcome of epilepsy in patients with Prader-Willi syndrome*. *J Neurol*. 2015 Jan;262(1):116-23. doi: 10.1007/s00415-014-7542-1. Epub 2014 Oct 18. Erratum in: *J Neurol*. 2015 Jan;262(1):124-5. Carotenuto, Marco [added]; Esposito, Maria [added]. PubMed PMID: 25326049.
- 32) Iber, C., Ancoli-Israel, S., Chesson, A., Quan, S.F. *The AASM manual for the scoring of sleep and associated events: rules, terminology and technical specifications*. American Academy of Sleep Medicine, Westchester, IL; 2007.
- 33) *American Psychiatric Association, Diagnostic and statistical manual of mental disorders: DSM-IV-TR*, American Psychiatric Pub, 4th ed: 2000. ISBN 0-89042-025-4; ISBN 978-0-89042-025-6
- 34) Conners CK. 2000. *Conners' Rating Scales-Revised: Technical Manual*. North Tonawanda, NY:Multi Health Systems.
- 35) Chieffi S, Iavarone A, Iaccarino L, La Marra M, Messina G, De Luca V, Monda M. *Age-related differences in distractor interference on line bisection*. *Exp Brain Res*. 2014 Nov;232(11): 3659-64. doi: 10.1007/s00221-014-4056-0
- 36) Youssef NA, Ege M, Angly SS, Strauss JL, Marx CE. *Is obstructive sleep apnea associated with ADHD?* *Ann Clin Psychiatry*. 2011 Aug; 23(3): 213-24.
- 37) Geladé K, Janssen TW, Bink M, van Mourik R, Maras A, Oosterlaan J. *Behavioral Effects of Neurofeedback Compared to Stimulants and Physical Activity in Attention-Deficit/Hyperactivity Disorder: A Randomized Controlled Trial*. *J Clin Psychiatry*. 2016 Sep 13. doi: 10.4088/JCP.15m10149
- 38) Messina G, Di Bernardo G, Viggiano A, De Luca V, Monda V, Messina A, Chieffi S, Galderisi U, Monda M. *Exercise increases the level of plasma orexin A in humans*. *J Basic Clin Physiol Pharmacol*. 2016 Sep 26. pii: j/bcphp.ahead-of-print/jbcpp-2015-0133/jbcpp-2015-0133.xml. doi: 10.1515/jbcpp-2015-0133.
- 39) Messina G, Dalia C, Tafuri D, Monda V, Palmieri F, Dato A, Russo A, De Blasio S, Messina A, De Luca V, Chieffi S, Monda M. *Orexin-A controls sympathetic activity and eating behavior*. *Front Psychol*. 2014 Sep 8; 5: 997. doi:10.3389/fpsyg.2014.00997. eCollection 2014. Review.
- 40) Visser SN, Bitsko RH, Danielson ML, Ghandour RM, Blumberg SJ, Schieve LA, Holbrook JR, Wolraich ML, Cuffe SP. *Treatment of Attention Deficit/Hyperactivity Disorder among Children with Special Health Care Needs*. *J Pediatr*. 2015 Jun;166(6): 1423-30.e1-2. doi: 10.1016/j.jpeds.2015.02.018
- 41) Parisi, L Di Filippo T, Roccella M. *The child with autism spectrum disorders (ASD): behavioral and neurobiological aspects*. *Acta Medica Mediterranea*, 2015, 31: 1187.
- 42) Entesar Foumany GhH, Salehi J. *The relationship between emotional intelligence and life satisfaction and the mediatory role of resiliency and emotional balance among the students of zanjan university*. *Acta Medica Mediterranea*, 2015, 31: 1351.
- 43) Nabi Amjad R, Navab E, Nikbakht Nasrabad A. *Parents of children with epilepsy captured by epilepsy: a qualitative study*. *Acta Medica Mediterranea*, 2016, 32: 1303
- 44) Shams S, Omolbanin Mohammadian S, Monajemzadeh M, Emamgholipour S, Aghi Haghi Ashtiani M, Irani H, Shafeghat L. *Evaluation of serum transferrin receptors in children with iron deficiency anemia*. *Acta Medica Mediterranea*, 2016, 32: 1555.
- 45) Yongli Z, Chunting L. *Problems and countermeasures of pediatric emergency nursing security*. *Acta Medica Mediterranea*, 2016, 32: 1177.
- 46) Mirzaei-Alavijeh M, Hidarnia A, Kok G, Niknami S , Esmail Motlagh M. *Family-based cognitive factors effective on preventing the onset of substance use in iranian society's children: applying the intervention mapping protocol*. *Acta Medica Mediterranea*, 2016, 32: 1015.
- 47) Jalili Z , Jalili C , Sakhaie S. *Frequency of congenital heart diseases and the probable predisposing risk factors in children admitted to imam ali hospital, kerman-shah, iran, 2002-2012*. *Acta Medica Mediterranea*, 2016, 32: 915.
- 48) Katirci Y, Ocak T, Karamercan MA, Kocaşaban D, Serkan Yurdakul M, Başpınar I, Coşkun F. *Compliance with catch rules in administering computerized tomography scans to children admitted to the emergency department with minor head trauma*. *Acta Medica Mediterranea*, 2013, 29: 717.
- 49) Ruberto M, Precenzano F, Parisi L, Salerno M, Maltese A, Messina G, Roccella M. *Visuomotor integration skills in children affected by obstructive sleep apnea syndrome: a case-control study*. *Acta Medica Mediterranea*. 2016, 32: 1659; DOI: 10.19193/0393-6384_2016_5_146
- 50) Parisi L, Ruberto M, Precenzano F, Di Filippo T, Russotto C, Maltese A, Salerno M, Roccella M. *The quality of life in children with cerebral palsy*. *Acta Medica Mediterranea*, 2016, 32: 1665; DOI: 10.19193/0393-6384_2016_5_147

Corresponding author

MICHELE ROCCELLA; MD; PhD
 Department of Psychological
 Pedagogical and Educational Sciences
 University of Palermo
 (Italy)